

Infinitely many local minima of sequentially weakly lower semicontinuous functionals and applications

BIAGIO RICCERI

I am here concerned with an overview of some applications of Theorem 1 below that have been obtained in the last two years.

THEOREM 1 ([14]). - *Let X be a non-empty sequentially weakly closed set in a reflexive real Banach space, and let $\Phi, \Psi : X \rightarrow]-\infty, +\infty]$ be two sequentially weakly lower semicontinuous functionals. Assume also that Ψ is (strongly) continuous. Denote by I the set of all real numbers $\rho > \inf_X \Psi$ such that $\Psi^{-1}(]-\infty, \rho[)$ is bounded and intersects the domain of Φ . Assume that $I \neq \emptyset$. For each $\rho \in I$, put*

$$\varphi(\rho) = \inf_{x \in \Psi^{-1}(]-\infty, \rho[)} \frac{\Phi(x) - \inf_{(\Psi^{-1}(]-\infty, \rho[))_w} \Phi}{\rho - \Psi(x)},$$

where $(\Psi^{-1}(]-\infty, \rho[))_w$ is the closure of $\Psi^{-1}(]-\infty, \rho[)$ in the relative weak topology of X . Furthermore, set

$$\gamma = \liminf_{\rho \rightarrow (\sup I)^-} \varphi(\rho)$$

and

$$\delta = \liminf_{\rho \rightarrow \left(\inf_X \Psi\right)^+} \varphi(\rho).$$

Then, the following conclusions hold:

- (a) For each $\rho \in I$ and each $\mu > \varphi(\rho)$, the restriction of the functional $\Phi + \mu\Psi$ to $\Psi^{-1}(]-\infty, \rho[)$ has a global minimum.
- (b) If $\gamma < +\infty$, then, for each $\mu > \gamma$, the following alternative holds: either the restriction of $\Phi + \mu\Psi$ to $\Psi^{-1}(]-\infty, \sup I[)$ has a global minimum, or there exists a sequence $\{x_n\}$ of local minima of $\Phi + \mu\Psi$ such that $\Psi(x_n) < \sup I$ for all $n \in \mathbf{N}$ and $\lim_{n \rightarrow \infty} \Psi(x_n) = \sup I$.
- (c) If $\delta < +\infty$, then, for each $\mu > \delta$, there exists a sequence $\{x_n\}$ of local minima of $\Phi + \mu\Psi$, with $\lim_{n \rightarrow \infty} \Psi(x_n) = \inf_X \Psi$, which weakly converges to a global minimum of Ψ .

A first consequence of conclusion (a) of Theorem 1 is as follows:

THEOREM 2 ([14]). - *Let E be a reflexive real Banach space, X a closed, convex, unbounded subset of E , and $\Phi, \Psi : X \rightarrow \mathbf{R}$ two convex functionals, with Φ lower semicontinuous and Ψ continuous and satisfying $\lim_{x \in X, \|x\| \rightarrow +\infty} \Psi(x) = +\infty$. Put*

$$\lambda^* = \inf_{\rho > \inf_X \Psi} \inf_{x \in \Psi^{-1}(]-\infty, \rho[)} \frac{\Phi(x) - \inf_{\Psi^{-1}(]-\infty, \rho[)} \Phi}{\rho - \Psi(x)}.$$

Then, for each $\lambda > \lambda^*$, the functional $\Phi + \lambda\Psi$ has a global minimum in X . Moreover, if $\lambda^* > 0$, for each $\mu < \lambda^*$, the functional $\Phi + \mu\Psi$ has no global minima in X .

REMARK 1. - The second conclusion of Theorem 2 does not hold, in general, if $\lambda^* = 0$. To see this, consider, for instance, the case when $\Phi(x) = \|x\|^2$ and $\Psi(x) = \|x\|$.

Let us recall that a Gâteaux differentiable functional J on a real Banach space X is said to satisfy the Palais-Smale condition if each sequence $\{x_n\}$ in X such that $\sup_{n \in \mathbf{N}} |J(x_n)| < +\infty$ and $\lim_{n \rightarrow +\infty} \|J'(x_n)\|_{X^*} = 0$ admits a strongly converging subsequence.

A joint application of conclusion (a) of Theorem 1 with the main result of [13] gives the following two critical points theorem:

THEOREM 3 ([17]). - Let X be a reflexive real Banach spaces and let $\Phi, \Psi : X \rightarrow \mathbf{R}$ be two sequentially weakly lower semicontinuous and continuously Gâteaux differentiable functionals. In addition, assume that, for each $\lambda > 0$, the functional $\Phi + \lambda\Psi$ satisfies the Palais-Smale condition.

Then, for each $\rho > \inf_X \Psi$ and each μ satisfying

$$\mu > \inf_{x \in \Psi^{-1}(]-\infty, \rho[)} \frac{\Phi(x) - \inf_{(\Psi^{-1}(]-\infty, \rho[))_w} \Phi}{\rho - \Psi(x)} ,$$

the following alternative holds: either the functional $\Phi + \mu\Psi$ has a strict global minimum which lies in $\Psi^{-1}(]-\infty, \rho[)$, or the same functional has at least two critical points one of which lies in $\Psi^{-1}(]-\infty, \rho[)$.

We also recall that if X is a real Hilbert space, an operator $A : X \rightarrow X$ is said to be a potential operator if there exists a Gâteaux differentiable functional P on X (which is called a potential of A) such that $P' = A$.

Applying Theorem 1, we get the following result about fixed points of potential operators in real Hilbert spaces:

THEOREM 4 ([14]). - Let X be a real Hilbert space, and let $A : X \rightarrow X$ be a potential operator, with sequentially weakly upper semicontinuous potential P . For each $\rho > 0$, put

$$\varphi(\rho) = \inf_{\|x\|^2 < \rho} \frac{\sup_{\|y\|^2 \leq \rho} P(y) - P(x)}{\rho - \|x\|^2} .$$

Furthermore, set

$$\gamma = \liminf_{\rho \rightarrow +\infty} \varphi(\rho)$$

and

$$\delta = \liminf_{\rho \rightarrow 0^+} \varphi(\rho) .$$

Then, the following conclusions hold:

- (a) If there is $\rho > 0$ such that $\varphi(\rho) < \frac{1}{2}$, then the operator A has a fixed point whose norm is less than $\sqrt{\rho}$.
- (b) If $\gamma < \frac{1}{2}$, then the following alternative holds: either the functional $x \rightarrow \frac{1}{2}\|x\|^2 - P(x)$ has a global minimum, or the set of all fixed points of A is unbounded.
- (c) If $\delta < \frac{1}{2}$, then the following alternative holds: either 0 is a local minimum of the functional $x \rightarrow \frac{1}{2}\|x\|^2 - P(x)$, or there exists a sequence of pairwise distinct fixed points of A which strongly converges to 0.

In particular, we have

THEOREM 5 ([14]). - Let X be a real Hilbert space, and let $A : X \rightarrow X$ be a potential operator, with sequentially weakly upper semicontinuous potential P satisfying

$$\liminf_{r \rightarrow +\infty} \frac{\sup_{\|x\| \leq r} P(x)}{r^2} < \frac{1}{2} < \limsup_{r \rightarrow +\infty} \frac{\sup_{\|x\| \leq r} P(x)}{r^2}.$$

Then, the set of all fixed points of A is unbounded.

From now on, Ω is an open bounded subset of $\mathbf{R}^n$, with smooth boundary, and (for $p > 1$) $W^{1,p}(\Omega)$, $W_0^{1,p}(\Omega)$ are the usual Sobolev spaces, with norms

$$\|u\| = \left(\int_{\Omega} |\nabla u(x)|^p dx + \int_{\Omega} |u(x)|^p dx \right)^{\frac{1}{p}}$$

and

$$\|u\| = \left(\int_{\Omega} |u(x)|^p dx \right)^{\frac{1}{p}}$$

respectively.

Let $p > 1$, and let $\phi : \Omega \times \mathbf{R} \rightarrow \mathbf{R}$ be a Carathéodory function.

Recall that a weak solution of the Dirichlet problem

$$\begin{cases} -\operatorname{div}(|\nabla u|^{p-2} \nabla u) = f(x, u) & \text{in } \Omega \\ u|_{\partial\Omega} = 0 \end{cases}$$

is any $u \in W_0^{1,p}(\Omega)$ such that

$$\int_{\Omega} |\nabla u(x)|^{p-2} \nabla u(x) \nabla v(x) dx - \int_{\Omega} f(x, u(x)) v(x) dx = 0 \quad (*)$$

for all $v \in W_0^{1,p}(\Omega)$. While, a weak solution of the Neumann problem

$$\begin{cases} -\operatorname{div}(|\nabla u|^{p-2} \nabla u) = f(x, u) & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega \end{cases}$$

ν being the outer unit normal to $\partial\Omega$, is is any $u \in W^{1,p}(\Omega)$ satisfying identity (*) for all $v \in W^{1,p}(\Omega)$.

The following is one of the most classical existence results on the Dirichlet problem for nonlinear elliptic equations:

THEOREM A ([1]). - *Assume that:*

(1) *there are two positive constants a, q , with $q < \frac{n+2}{n-2}$ if $n \geq 3$, such that*

$$|f(x, \xi)| \leq a(1 + |\xi|^q)$$

for all $(x, \xi) \in \Omega \times \mathbf{R}$;

(2) *there are constants $r \geq 0$ and $c > 2$ such that*

$$0 < c \int_0^\xi f(x, t) dt \leq \xi f(x, \xi)$$

for all $(x, \xi) \in \Omega \times \mathbf{R}$ with $|\xi| \geq r$;

(3) *one has*

$$\lim_{\xi \rightarrow 0} \frac{f(x, \xi)}{\xi} = 0$$

uniformly with respect to x .

Then, the problem

$$\begin{cases} -\Delta u = f(x, u) & \text{in } \Omega \\ u|_{\partial\Omega} = 0. \end{cases}$$

has a non zero weak solution.

What can be said whether, in Theorem A, condition (3) is removed at all ? Using Theorem 3, one obtains the following

THEOREM 6 ([17]). - *Assume that conditions (1) and (2) hold.*

Then, for each $\rho > 0$ and each μ satisfying

$$\mu > \inf_{u \in B_\rho} \frac{\sup_{v \in B_\rho} \int_\Omega \left(\int_0^{v(x)} f(x, \xi) d\xi \right) dx - \int_\Omega \left(\int_0^{u(x)} f(x, \xi) d\xi \right) dx}{\rho - \int_\Omega |\nabla u(x)|^2 dx}, \quad (**)$$

where

$$B_\rho = \left\{ u \in W_0^{1,2}(\Omega) : \int_\Omega |\nabla u(x)|^2 dx < \rho \right\},$$

the problem

$$\begin{cases} -\Delta u = \frac{1}{2\mu} f(x, u) & \text{in } \Omega \\ u|_{\partial\Omega} = 0 \end{cases}$$

has at least two weak solutions one of which lies in B_ρ .

The following problem naturally arises in connection with Theorems A and 6.

PROBLEM 1. - Under conditions (1) and (2) of Theorem A, is there some $\rho > 0$ such that the infimum appearing in (**) is less than $\frac{1}{2}$?

Clearly, if the answer to this problem was positive, then Theorem 6 would be a proper improvement of Theorem A.

Another result proved using Theorem 1 is the following:

THEOREM 7 ([18]). - Let $f, g : \Omega \times \mathbf{R} \rightarrow \mathbf{R}$ be two Carathéodory functions. Assume that:

(i) there is $s > 1$ such that

$$\limsup_{\xi \rightarrow 0^+} \frac{\sup_{x \in \Omega} |f(x, \xi)|}{\xi^s} < +\infty ;$$

(ii) there is $q \in]0, 1[$ such that

$$\limsup_{\xi \rightarrow 0^+} \frac{\sup_{x \in \Omega} |g(x, \xi)|}{\xi^q} < +\infty ;$$

(iii) there are a non-empty open set $D \subseteq \Omega$ and a set $B \subseteq D$ of positive measure such that

$$\limsup_{\xi \rightarrow 0^+} \frac{\inf_{x \in B} \int_0^\xi g(x, t) dt}{\xi^2} = +\infty , \quad \liminf_{\xi \rightarrow 0^+} \frac{\inf_{x \in D} \int_0^\xi g(x, t) dt}{\xi^2} > -\infty .$$

Then, for some $\lambda^* > 0$ and for each $\lambda \in]0, \lambda^*[$, the problem

$$\begin{cases} -\Delta u = f(x, u) + \lambda g(x, u) & \text{in } \Omega \\ u|_{\partial\Omega} = 0 , \end{cases} \quad (P_\lambda)$$

admits a non-zero, non-negative weak solution $u_\lambda \in C^1(\overline{\Omega})$. Moreover, one has

$$\limsup_{\lambda \rightarrow 0^+} \frac{\|u_\lambda\|_{C^1(\overline{\Omega})}}{\lambda^{\frac{q}{1-q}}} < +\infty$$

and the function

$$\lambda \rightarrow \frac{1}{2} \int_{\Omega} |\nabla u_\lambda(x)|^2 dx - \int_{\Omega} \left(\int_0^{u_\lambda(x)} f(x, \xi) d\xi \right) dx - \lambda \int_{\Omega} \left(\int_0^{u_\lambda(x)} g(x, \xi) d\xi \right) dx$$

is negative and decreasing in $]0, \lambda^*[$. If, in addition, f, g are continuous in $\Omega \times [0, +\infty[$ and

$$\liminf_{\xi \rightarrow 0^+} \frac{\inf_{x \in \Omega} g(x, \xi)}{\xi |\log \xi|^2} > -\infty ,$$

then u_λ is positive in Ω .

REMARK 2. - Observe that Theorem 7 is bifurcation result. In fact, it ensures, in particular, that $\lambda = 0$ is a bifurcation point for problem (P_λ) , in the sense that $(0, 0)$ belongs to the closure in $C^1(\overline{\Omega}) \times \mathbf{R}$ of the set

$$\{(u, \lambda) \in C^1(\overline{\Omega}) \times]0, +\infty[: u \text{ is a weak solution of } (P_\lambda), u \neq 0, u \geq 0\} .$$

In view of [2], where problem (P_λ) is studied for particular nonlinearities, the following problems arises:

PROBLEM 2. - Under the assumptions of Theorem 7, does problem (P_λ) admit a non-zero, non-negative, *minimal* solution for each $\lambda > 0$ small enough ?

REMARK 3. - Again applying Theorem 1, some bifurcations theorem for Hammerstein nonlinear integral equations (in the spirit of Theorem 7) have been obtained by F. Faraci in [10].

Note, in particular, the following corollary of Theorem 7:

COROLLARY 1 ([18]). - Let $0 < q < 1 < s$ and let α, β be two Hölder continuous functions on $\overline{\Omega}$. Assume that

$$0 \leq \inf_{\Omega} \beta, \quad 0 < \sup_{\Omega} \beta .$$

Then, for some $\lambda^* > 0$ and for each $\lambda \in]0, \lambda^*[$, the problem

$$\begin{cases} -\Delta u = \alpha(x)u^s + \lambda\beta(x)u^q & \text{in } \Omega \\ u|_{\partial\Omega} = 0 \end{cases}$$

admits a positive classical solution u_λ . Moreover, one has

$$\limsup_{\lambda \rightarrow 0^+} \frac{\|u_\lambda\|_{C^1(\overline{\Omega})}}{\lambda^{\frac{q}{1-q}}} < +\infty$$

and the function

$$\lambda \rightarrow \frac{1}{2} \int_{\Omega} |\nabla u_\lambda(x)|^2 dx - \frac{1}{s+1} \int_{\Omega} \alpha(x) |u_\lambda(x)|^{s+1} dx - \frac{\lambda}{q+1} \int_{\Omega} \beta(x) |u_\lambda(x)|^{q+1} dx$$

is negative and decreasing in $]0, \lambda^*[$.

In the next theorems, λ denotes a function in $L^\infty(\Omega)$, with $\text{ess inf}_\Omega \lambda > 0$. They have been obtained in [16] as applications of conclusions (b) and (c) of Theorem 1.

THEOREM 8 ([16]). - Assume $p > n$. Let $f : \mathbf{R} \rightarrow \mathbf{R}$ be a continuous function, and $\{a_k\}, \{b_k\}$ two sequences in $\mathbf{R}^+$ satisfying

$$a_k < b_k \quad \forall k \in \mathbf{N}, \quad \lim_{k \rightarrow \infty} b_k = +\infty, \quad \lim_{k \rightarrow \infty} \frac{a_k}{b_k} = 0 ,$$

$$\max \left\{ \sup_{\xi \in [a_k, b_k]} \int_{a_k}^{\xi} f(t) dt, \sup_{\xi \in [-b_k, -a_k]} \int_{-a_k}^{\xi} f(t) dt \right\} \leq 0 \quad \forall k \in \mathbf{N}$$

and

$$\limsup_{|\xi| \rightarrow +\infty} \frac{\int_0^{\xi} f(t) dt}{|\xi|^p} = +\infty .$$

Then, for every $\alpha, \beta \in L^1(\Omega)$, with $\min\{\alpha(x), \beta(x)\} \geq 0$ a.e. in Ω and $\alpha \neq 0$, and for every continuous function $g : \mathbf{R} \rightarrow \mathbf{R}$ satisfying

$$\sup_{\xi \in \mathbf{R}} \int_0^{\xi} g(t) dt \leq 0$$

and

$$\liminf_{|\xi| \rightarrow +\infty} \frac{\int_0^{\xi} g(t) dt}{|\xi|^p} > -\infty ,$$

the problem

$$\begin{cases} -\text{div}(|\nabla u|^{p-2} \nabla u) + \lambda(x)|u|^{p-2}u = \alpha(x)f(u) + \beta(x)g(u) & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega \end{cases}$$

admits an unbounded sequence of weak solutions in $W^{1,p}(\Omega)$.

A typical example of application of Theorem 8 is as follows:

EXAMPLE 1 ([16]). - Let $p > n$. Then, for each $\eta \in L^1(\Omega)$ with $\text{ess inf}_\Omega \eta > 0$, the problem

$$\begin{cases} -\text{div}(|\nabla u|^{p-2} \nabla u) = \eta(x) \left(\sum_{k=1}^{\infty} (\text{dist}(u, \mathbf{R} \setminus [k!k, (k+1)!])^p - |u|^{p-2}u) \right) & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega \end{cases}$$

admits an unbounded sequence of weak solutions in $W^{1,p}(\Omega)$.

THEOREM 9 ([16]). - Assume $p > n$. Let $f : \mathbf{R} \rightarrow \mathbf{R}$ be a continuous function, and $\{a_k\}, \{b_k\}$ two sequences in $\mathbf{R}^+$ satisfying

$$a_k < b_k \quad \forall k \in \mathbf{N}, \quad \lim_{k \rightarrow \infty} b_k = 0, \quad \lim_{k \rightarrow \infty} \frac{a_k}{b_k} = 0 ,$$

$$\max \left\{ \sup_{\xi \in [a_k, b_k]} \int_{a_k}^{\xi} f(t) dt, \sup_{\xi \in [-b_k, -a_k]} \int_{-a_k}^{\xi} f(t) dt \right\} \leq 0 \quad \forall k \in \mathbf{N}$$

and

$$\limsup_{\xi \rightarrow 0} \frac{\int_0^{\xi} f(t) dt}{|\xi|^p} = +\infty .$$

Then, for every $\alpha, \beta \in L^1(\Omega)$, with $\min\{\alpha(x), \beta(x)\} \geq 0$ a.e. in Ω and $\alpha \neq 0$, and for every continuous function $g : \mathbf{R} \rightarrow \mathbf{R}$ satisfying

$$\sup_{\xi \in \mathbf{R}} \int_0^{\xi} g(t) dt \leq 0$$

and

$$\liminf_{\xi \rightarrow 0} \frac{\int_0^{\xi} g(t) dt}{|\xi|^p} > -\infty ,$$

the problem

$$\begin{cases} -\operatorname{div}(|\nabla u|^{p-2} \nabla u) + \lambda(x)|u|^{p-2}u = \alpha(x)f(u) + \beta(x)g(u) & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega \end{cases}$$

admits a sequence of non zero weak solutions which strongly converges to 0 in $W^{1,p}(\Omega)$.

REMARK 4. - Again using Theorem 1, Theorems 8 and 9 have been extended to the setting of elliptic variational-hemivariational inequalities by D. Motreanu and S. A. Marano in [12]. While P. Candito, in [8], extended them to the case of discontinuous nonlinearities. Further various applications of Theorem 1 can be found in [3], [4], [5], [9], [11] and [15]. See also the related papers [6] and [19].

PROBLEM 3. - In Theorems 8 and 9, when $g = 0$, are the conclusions still valid without the assumption

$$\lim_{k \rightarrow \infty} \frac{a_k}{b_k} = 0 ?$$

A partial answer to this problem has recently been provided by G. Anello and G. Cordaro in [7].

References

- [1] A. AMBROSETTI and P. H. RABINOWITZ, *Dual variational methods in critical point theory and applications*, J. Funct. Anal., **14** (1973), 349-381.
- [2] A. AMBROSETTI, H. BREZIS and G. CERAMI, *Combined effects of concave and convex nonlinearities in some elliptic problems*, J. Funct. Anal., **122** (1994), 519-543.
- [3] G. ANELLO and G. CORDARO, *Existence of solutions of the Neumann problem involving the p -Laplacian via a variational principle of Ricceri*, Arch. Math. (Basel), to appear.
- [4] G. ANELLO and G. CORDARO, *An existence theorem for the Neumann problem involving the p -Laplacian*, preprint.
- [5] G. ANELLO and G. CORDARO, *An existence and localization theorem of solution for a Dirichlet problem*, preprint.
- [6] G. ANELLO and G. CORDARO, *Positive infinitely many and arbitrarily small solutions for the Dirichlet problem involving the p -Laplacian*, Proc. Royal Soc. Edinburgh Sect. A, to appear.
- [7] G. ANELLO and G. CORDARO, *Infinitely many positive solutions for the Neumann problem involving the p -Laplacian*, preprint.
- [8] P. CANDITO, *Infinitely many solutions to the Neumann problem for elliptic equations involving the p -Laplacian and with discontinuous nonlinearities*, Proc. Math. Soc. Edinburgh, to appear.
- [9] G. CORDARO, *Existence and location of periodic solutions to convex and non coercive Hamiltonian systems*, preprint.
- [10] F. FARACI, *Bifurcation theorems for Hammerstein nonlinear integral equations*, Glasgow Math. J., to appear.
- [11] F. FARACI, *Multiplicity results for a Neumann problem involving the p -Laplacian*, preprint.
- [12] S. A. MARANO and D. MOTREANU, *Infinitely many critical points of non-differentiable functions and applications to a Neumann type problem involving the p -Laplacian*, J. Differential Equations, to appear.
- [13] P. PUCCI and J. SERRIN, *A mountain pass theorem*, J. Differential Equations, **60** (1985), 142-149.
- [14] B. RICCERI, *A general variational principle and some of its applications*, J. Comput. Appl. Math., **113** (2000), 401-410.
- [15] B. RICCERI, *Existence and location of solutions to the Dirichlet problem for a class of nonlinear elliptic equations*, Appl. Math. Lett., **14** (2001), 143-148.
- [16] B. RICCERI, *Infinitely many solutions of the Neumann problem for elliptic equations involving the p -Laplacian*, Bull. London Math. Soc., **33** (2001), 331-340.
- [17] B. RICCERI, *On a classical existence theorem for nonlinear elliptic equations*, in "Experimental, constructive and nonlinear analysis", M. Théra ed., 275-278, CMS Conf. Proc. **27**, Canad. Math. Soc., 2000.

- [18] B. RICCERI, *A bifurcation theorem for nonlinear elliptic equations*, preprint.
- [19] J. SAINT RAYMOND, *On the multiplicity of the solutions of the equation $-\Delta u = \lambda f(u)$* , J. Differential Equations, to appear.

Department of Mathematics
University of Catania
Viale A. Doria 6
95125 Catania, Italy